



Laboratory Guidebook Notice of Change

Chapter new, **revised**, or archived: MLG 4.10

Title: Isolation and Identification of *Salmonella* from Meat, Poultry, Pasteurized Egg, and Siluriformes (Fish) Products and Carcass and Environmental Sponges

Effective Date: 1/02/19

Description and purpose of change(s):

The rapid molecular screening assay was revised in the equipment section and additional rapid screen instructions were added to the procedure.

Whole genome sequencing was included as an option to further characterize isolates during confirmation in lieu of serological testing with Somatic (O) Antigen and Flagellar (H) Antigen Agglutination Tests.

Additional instructions were added to perform a fluorescence check on the SBA plate during biochemical confirmation for the positive UV *Salmonella* control and presumptive positive samples following incubation.

Optional procedures not used by FSIS laboratories were removed from the method including references to XLT4 plating media and serology.

MLG 4C “FSIS Procedure for the Use of a Polymerase Chain Reaction (PCR) Assay for Screening *Salmonella* in Meat, Poultry, Egg, and Siluriformes (Fish) Products and Carcass and Environmental Sponges” will be archived as the rapid molecular screen procedure is now included in MLG 4.

**United States Department of Agriculture
Food Safety And Inspection Service, Office of Public Health Science**

MLG 4.10		Page 1 of 17
Title: Isolation And Identification of <i>Salmonella</i> From Meat, Poultry, Pasteurized Egg, and Siluriformes (Fish) Products and Carcass and Environmental Sponges		
Revision: .10	Replaces: .09	Effective: 01/02/19

Procedure Outline

- 4.1 Introduction
- 4.2 Safety Precautions
- 4.3 Quality Control Procedures
 - 4.3.1 Method Controls
 - 4.3.2 Specific Procedure Controls
- 4.4 Equipment, Reagents, Media and Test Kits
 - 4.4.1 Equipment
 - 4.4.2 Reagents and Test Systems
 - 4.4.3 Media
- 4.5 Sample Preparation
 - 4.5.1 Ready-to-Eat Meat, Poultry and Siluriformes Foods
 - 4.5.2 Raw Poultry Products
 - 4.5.3 Raw Meat and Raw Beef Mixed Products
 - 4.5.4 Carcass Sponges and Environmental Sponges
 - 4.5.5 Whole Bird and Parts Rinses
 - 4.5.6 Pasteurized Liquid, Frozen, or Dried Egg Products
 - 4.5.7 Raw Siluriformes (Fish) Products
 - 4.5.8 Fermented Products
 - 4.5.9 Dried Products (Breeding Mix, Dehydrated Sauce, Dried Soup Mix, and Dried Milk)
 - 4.5.10 Most Probable Numbers (MPN) Determination
- 4.6 Rapid Screening Salmonella Test Procedure
- 4.7 Selective Enrichment and Plating Media
- 4.8 Examination of and Picking Colonies from Plating Media
 - 4.8.1 Picking Colonies
 - 4.8.2 Screening Media
- 4.9 Biochemical Procedures
- 4.10 Culture Storage and Maintenance
- 4.11 Selected References

**United States Department of Agriculture
Food Safety And Inspection Service, Office of Public Health Science**

MLG 4.10		Page 2 of 17
Title: Isolation And Identification of <i>Salmonella</i> From Meat, Poultry, Pasteurized Egg, and Siluriformes (Fish) Products and Carcass and Environmental Sponges		
Revision: .10	Replaces: .09	Effective: 01/02/19

4.1 Introduction

The methods described in this guidebook are for use by the FSIS laboratories. FSIS does not specifically endorse any of the mentioned test products and acknowledges that equivalent products may be available for laboratory use. FSIS utilizes the following performance criteria when evaluating the suitability of an alternative laboratory method or product for a given analyte and sample matrix pair:

- Sensitivity of 90% or greater
- Specificity of 90% or greater
- Accuracy of 90% or greater
- Positive predictive value of 90% or greater
- Negative predictive value of 90% or greater

Performance criteria are relative to the reference cultural method for that analyte and sample matrix as outlined in the corresponding MLG chapter. Method validation is necessary to demonstrate the equivalence of alternative tests as detailed in the document titled “FSIS Guidance for Evaluating Test Kit Performance.”

This method describes the analysis of various meat, poultry and Siluriformes (fish) products, sponge and rinse samples, and egg products for *Salmonella*. It is not intended for the isolation and identification of *Salmonella typhi*.

Success in isolating *Salmonella* from any food can be related to a number of factors including food preparation procedures, the number of organisms present, sample handling after collection, etc. With raw samples, the competitive flora may be the most important factor. It varies from sample to sample and from one kind of matrix to another.

Another consideration is whether the examination is for routine monitoring or epidemiological purposes. The analyst may choose to augment the method for epidemiological purposes with additional enrichment procedures and culture media, two temperatures of incubation, intensified selection of colonies from plates, and/or rapid screening methods.

Unless otherwise stated all measurements cited in this method have a tolerance range of $\pm 2\%$.

**United States Department of Agriculture
Food Safety And Inspection Service, Office of Public Health Science**

MLG 4.10	Page 3 of 17	
Title: Isolation And Identification of <i>Salmonella</i> From Meat, Poultry, Pasteurized Egg, and Siluriformes (Fish) Products and Carcass and Environmental Sponges		
Revision: .10	Replaces: .09	Effective: 01/02/19

4.2 Safety Precautions

Salmonella are generally categorized as Biosafety Level 2 pathogens. CDC guidelines for handling Biosafety Level 2 pathogens should be followed whenever live cultures of *Salmonella* are used. A Class II laminar flow biosafety cabinet is recommended for procedures in which infectious aerosols or splashes may be created. The Safety Data Sheet (SDS) must be obtained from the manufacturer for the media, chemicals, reagents and microorganisms used in the analysis. The personnel who will handle the material should read the SDS prior to startup.

4.3 Quality Control Procedures

4.3.1 Method Controls

A *Salmonella* sp. H₂S positive culture and an uninoculated media control must be used from the start of the analysis. A H₂S negative *Salmonella* sp. positive culture shall also be included either from the start of the analysis or limited to sets containing screen positive samples, from the differentiation and confirmation steps starting with streaking to BGS and DMLIA agar plates. To facilitate identification of control isolates, the laboratory may use strains of uncommonly found serogroups or tagged cultures such as those that visibly fluoresce under ultraviolet (UV) light to differentiate QC strains from true contaminants. *S. Abaetetuba* is suggested as a readily available, H₂S positive culture that is not commonly found in meats or meat products. *S. Choleraesuis* is typically negative for H₂S production. These cultures may be obtained from ATCC. Other serotypes may be found that have H₂S negative strains. The positive control cultures should be inoculated into an appropriate matrix at a low inoculum level, e.g., by preparing a test organism suspension in broth or saline equivalent in turbidity to a 0.5 McFarland standard. Using a 1µL loop, inoculate the broth or streak the plates to be tested. Alternatively, commercially prepared bacterial pellets may be used. Once the control cultures are started, incubate the controls along with the samples, and analyze them in the same manner as the samples. Confirm at least one isolate from the H₂S positive control sample. Confirmation of at least one colony from the H₂S negative control is required when confirming H₂S negative samples. In the absence of a positive test sample, control cultures may be terminated at the same point as the sample analyses.

**United States Department of Agriculture
Food Safety And Inspection Service, Office of Public Health Science**

MLG 4.10	Page 4 of 17	
Title: Isolation And Identification of <i>Salmonella</i> From Meat, Poultry, Pasteurized Egg, and Siluriformes (Fish) Products and Carcass and Environmental Sponges		
Revision: .10	Replaces: .09	Effective: 01/02/19

4.3.2 Specific Procedure Controls

Each step of the analysis requires the use of appropriate controls to verify that the results are valid. Biochemical kit and rapid test manufacturers may specify control cultures for use with their products. If not specified, quality control procedures for biochemical tests and test media should include cultures that will demonstrate pertinent characteristics of the product.

4.4 Equipment, Reagents, Media and Test Kits

Not all of the materials listed below may be needed. Media and reagents specific to the selected biochemical test method may be needed in addition to the materials listed below.

4.4.1 Equipment

- a. Sterile tablespoons, scissors, forceps, knives, glass stirring rods, pipettes, petri dishes, test tubes, bent glass rods ("hockey sticks") as needed
- b. Blending/mixing equipment: Paddle blender, Sterile Osterizer-type blender with sterilized cutting assemblies, and blender jars or equivalent and adapters for use with Mason jars;
- c. Sterile plain, clear polypropylene bags (ca. 24" x 30 - 36"), or Whirl-pak type bags, or equivalent
- d. Incubator, $35 \pm 2^{\circ}\text{C}$
- e. Incubator or water bath, $42 \pm 0.5^{\circ}\text{C}$
- f. Balance, 2000 g capacity, sensitivity of 0.1 g
- g. Inoculating needles and loops
- h. Vortex mixer
- i. UV lamp, blue light excitation 475-495 nm light
- j. 3M™ Molecular Detection System
 - 3M™ Molecular Detection Heat Block Insert MDSHBIN
 - 3M™ Molecular Detection Speed Loader Tray MDSSLT
 - 3M™ Molecular Detection Cap/ Decap Tool-Lysis
 - 3M™ Molecular Detection Chill Block Insert MDSCBIN held at $20\text{-}25^{\circ}\text{C}$
 - VWR Digital 2 Block Heater or equivalent set at $100 \pm 1^{\circ}\text{C}$
 - Pipettor to deliver 20 μL and sterile disposable filter tips
 - Multichannel Pipettor to deliver 20 μL
- k. VITEK® 2 Compact System, or equivalent

**United States Department of Agriculture
Food Safety And Inspection Service, Office of Public Health Science**

MLG 4.10	Page 5 of 17	
Title: Isolation And Identification of <i>Salmonella</i> From Meat, Poultry, Pasteurized Egg, and Siluriformes (Fish) Products and Carcass and Environmental Sponges		
Revision: .10	Replaces: .09	Effective: 01/02/19

4.4.2 Reagents and Test Systems

- a. Crystal violet dye, 1% aqueous solution
- b. Saline, 0.85%
- c. Calcium carbonate, sterile
- d. 3M™ Molecular Detection Assay 2 - Salmonella
- e. Biochemical test panel (GN cards for VITEK® 2 Compact System)

4.4.3 Media

- a. Buffered peptone water (BPW) or Modified Tryptone Soya Broth (mTSB)
- b. TT broth (Hajna)
- c. Modified Rappaport Vassiliadis (mRV) broth, Rappaport-Vassiliadis R10 broth, or Rappaport-Vassiliadis Soya Peptone Broth (RVS)
- d. Brilliant green sulfa agar (BGS; contains 0.1% sodium sulfapyridine)
- e. Double modified lysine iron agar (DMLIA)
- f. Triple sugar iron agar (TSI)
- g. Lysine iron agar (LIA)
- h. Nutrient agar slants
- i. Tryptic soy agar with 5% sheep blood agar
- j. Additional media as needed for biochemical tests

4.5 Sample Preparation

Intact retail packages must be disinfected at the incision sites immediately prior to incision for sampling using an appropriate disinfectant, e.g., 3% hydrogen peroxide, ca. 70% ethanol or ca. 70% isopropanol. If the package does not appear to be clean, scrub gently using soapy water and rinse thoroughly prior to disinfection. A sterile scalpel may be helpful for cutting the packaging. Aseptically pull the packaging away to expose the product for sampling.

Note: For Ready-to-Eat (RTE) sausages in casing, the shell/casing is an integral part of the sample and should be free of pathogens and toxins. The casing is not to be disinfected since some casings are permeable and the disinfectant may be introduced into the core of the product. In addition consumers often slice through an inedible casing and then remove it thus any contamination on the surface of the casing could be transferred to the edible core of the product. Sample preparation and enrichment incubation times may vary by matrix and program. Refer to Table 1 and the following sample preparation sections.

**United States Department of Agriculture
Food Safety And Inspection Service, Office of Public Health Science**

MLG 4.10		Page 6 of 17
Title: Isolation And Identification of <i>Salmonella</i> From Meat, Poultry, Pasteurized Egg, and Siluriformes (Fish) Products and Carcass and Environmental Sponges		
Revision: .10	Replaces: .09	Effective: 01/02/19

Table 1. Sample Preparation and Enrichment Guide

Product	Sample Preparation		Incubation
	Portion Size	Enrichment Amount determined by volume or weight	Cultural or rapid screen
Ready-to-Eat Meat, Poultry and Siluriformes Foods	325 ± 6.5 g	975 ± 19.5 ml BPW	35 ± 2°C for 18-24 h
Raw Poultry Products	325 ± 32.5 g or 25 ± 2.5 g	1625 ± 32.5ml BPW or 225 ± 4.5 ml BPW	35 ± 2°C for 20-24 h
Raw Meat and Raw Beef Mixed Products	325 ± 32.5 g or 25 ± 2.5 g	975± 19.5 ml mTSB or 75 ± 1.5 ml mTSB	42 ± 1°C for 15-24 h
Poultry Carcass and Environmental Sponges	1 sponge pre-moistened with 10 ml buffer	50 ± 1 ml BPW to bring total volume to 60 ml*	35 ± 2°C for 20-24 h
Meat Carcass and Environmental Sponges	1 sponge pre-moistened with 10 ml buffer	50 ± 1 ml mTSB to bring total volume to 60 ml*	42 ± 1°C for 15-24 h
Whole Bird and Parts Rinses	30 ± 0.6 ml sample rinse fluid	30 ± 0.6 ml BPW	35 ± 2°C for 20-24 h
Pasteurized Liquid, Frozen or Dried Egg Products	100 ± 2 g	900 ± 18 ml BPW	35 ± 2°C for 18-24 h
Raw Siluriformes Products	25 ± 2.5 g	225 ± 4.5 ml BPW	35 ± 2°C for 22-26 h
Fermented Products	325 ± 6.5 g + 10 g of sterilized calcium carbonate	2925 ± 58.5 ml of BPW with 1 ml of a 1% aqueous solution of crystal violet per liter	35 ± 2°C for 18-24 h
Dried Products (Breeding Mix, Dehydrated Sauce, Soup Mix, Dried Milk)	325 ± 6.5 g	2925 ± 58.5 ml BPW	35 ± 2°C for 18-24 h

* or maintain a 1:6 ratio for different project buffer volumes, e.g., 25 ml buffer + 125 ml enrichment

**United States Department of Agriculture
Food Safety And Inspection Service, Office of Public Health Science**

MLG 4.10		Page 7 of 17
Title: Isolation And Identification of <i>Salmonella</i> From Meat, Poultry, Pasteurized Egg, and Siluriformes (Fish) Products and Carcass and Environmental Sponges		
Revision: .10	Replaces: .09	Effective: 01/02/19

4.5.1 Ready-to-Eat Meat, Poultry and Siluriformes Foods

Follow additional program requirements for preparing sample and sub-sample composites. Outbreak samples may require a different sample preparation. Follow customer specifications.

Using a sterile scalpel, knife, spoon, chisel or other tool, cut small pieces from representative sites of submitted product to prepare a composite sample portion. While multiple packages of a product are usually submitted, for large products a single package may be submitted.

For multi-component RTE products, follow the appropriate sample preparation instructions listed below:

If the meat, poultry or Siluriformes component is separate and distinct from other non-meat ingredients, analyze only the representative meat/poultry portion of the RTE product. Examples include products with the meat/poultry portion separate from any vegetable/dessert component, or fajita kits with meat/poultry, onions/peppers, and tortillas in three separate internal packages/bags within an outer package.

When meat, poultry, or Siluriformes is combined with other ingredients to form the product (e.g., beef stew containing vegetables, potatoes, etc.), analyze representative meat/poultry portions in combination with other ingredients.

- a. Weigh the composite sample into a large sterile bag (or sterile blender jar if required by the customer or sample type).
- b. Add ambient temperature sterile BPW. Blend or stomach approximately two minutes.
- c. Incubate at $35 \pm 2^{\circ}\text{C}$ for 18-24 h.
- d. Proceed to Section 4.6 for use of the rapid screen or refer to Section 4.7 to continue the cultural analysis.

**United States Department of Agriculture
Food Safety And Inspection Service, Office of Public Health Science**

MLG 4.10	Page 8 of 17	
Title: Isolation And Identification of <i>Salmonella</i> From Meat, Poultry, Pasteurized Egg, and Siluriformes (Fish) Products and Carcass and Environmental Sponges		
Revision: .10	Replaces: .09	Effective: 01/02/19

4.5.2 Raw Poultry Products

- a. Unless the exact sample portion is submitted, weigh the product into a sterile polypropylene bag, sterile blender jar, or other sterile jar. Note: If the sample is not already ground, in some cases it may be best to mince it with sterile scissors or leave it whole (e.g. chicken wings) to avoid jamming blender blades with skin or connective tissue.
- b. Add the BPW. Stomach, blend or hand massage until clumps are dispersed.
- c. Incubate at $35 \pm 2^{\circ}\text{C}$ for 20-24 h.
- d. Proceed to Section 4.6 for use of the rapid screen or refer to Section 4.7 to continue the cultural analysis.

4.5.3 Raw Meat and Raw Beef Mixed Products

- a. Unless the exact sample portion is submitted, weigh the meat into a sterile polypropylene bag, sterile blender jar, or other sterile jar.
- b. Add the mTSB at a 1:4 dilution, e.g., 325 ± 32.5 g sample with 975 ± 19.5 ml mTSB broth. Stomach, blend or hand massage until clumps are dispersed.
- c. Incubate at $42 \pm 1^{\circ}\text{C}$ for 15-24 h.
- d. Proceed to Section 4.6 for use of the rapid screen or refer to Section 4.7 to continue the cultural analysis.

4.5.4 Carcass Sponges and Environmental Sponges

- a. For poultry carcass or environmental sponges, add the BPW to the sample bag containing a sponge moistened with 10 ml of buffer to bring the total volume to 60 ml. Mix well. If the project calls for a larger volume of buffer, adjust the volume of enrichment broth to a 1:6 dilution. Incubate at $35 \pm 2^{\circ}\text{C}$ for 20-24 h.
- b. For meat carcass or environmental sponges, add the mTSB to the sample bag containing a sponge moistened with 10 ml of buffer to bring the total volume to

**United States Department of Agriculture
Food Safety And Inspection Service, Office of Public Health Science**

MLG 4.10		Page 9 of 17
Title: Isolation And Identification of <i>Salmonella</i> From Meat, Poultry, Pasteurized Egg, and Siluriformes (Fish) Products and Carcass and Environmental Sponges		
Revision: .10	Replaces: .09	Effective: 01/02/19

60 ml. Mix well. If the project calls for a larger volume of buffer, adjust the volume of enrichment broth to a 1:6 dilution. Incubate at $42 \pm 1^\circ\text{C}$ for 15-24 h.

- c. Proceed to Section 4.6 for use of the rapid screen or refer to Section 4.7 to continue the cultural analysis.

4.5.5 Whole Bird and Parts Rinses

Due to differences between sample types/sizes (e.g. chicken vs. turkey carcasses; size of parts), follow instructions given in the specific project protocol.

- a. For chicken carcasses:

Aseptically drain excess fluid from the carcass and transfer the carcass to a large sterile bag.

Pour 400 ml (or other volume specified in program protocol) of collection medium into the cavity of the carcass contained in the bag.

Rinse the bird inside and out with a rocking motion for one minute (ca. 35 RPM). This is done by grasping the carcass in the bag with one hand and the closed top of the bag with the other. Rock with a reciprocal motion in about an 18-24 inch arc, assuring that all surfaces (interior and exterior of the carcass) are rinsed.

- b. For chicken parts:

Add the weight of the specific part plus the volume of collection medium specified by the program protocol into the sterile bag.

Rinse the parts with the broth assuring that all surfaces are rinsed.

- c. Transfer the sample rinse fluid to a sterile container.
- d. Use 30 ± 0.6 ml of the sample rinse fluid obtained above for *Salmonella* analysis. Add 30 ± 0.6 ml of sterile BPW and mix well.
- e. Incubate at $35 \pm 2^\circ\text{C}$ for 20-24 h.

**United States Department of Agriculture
Food Safety And Inspection Service, Office of Public Health Science**

MLG 4.10	Page 10 of 17	
Title: Isolation And Identification of <i>Salmonella</i> From Meat, Poultry, Pasteurized Egg, and Siluriformes (Fish) Products and Carcass and Environmental Sponges		
Revision: .10	Replaces: .09	Effective: 01/02/19

- f. Proceed to Section 4.6 for use of the rapid screen or refer to Section 4.7 to continue the cultural analysis.

NOTE: If analyses other than *Salmonella* are to be performed, the carcass may be rinsed and dilutions made directly from the collection rinse.

4.5.6 Pasteurized Liquid, Frozen, or Dried Egg Products

- a. Mix the liquid sample with a sterile spoon, spatula, or by shaking.
- b. Weigh the liquid egg product into a sterile polypropylene bag, sterile blender jar or other sterile jar.
- c. Mix the inoculated BPW well by shaking, stomaching, or blending.

Note: If a special sample or specification requires a sample size other than 100 g, the dilution of egg sample to BPW is to be maintained at 1:10.

- d. With dried egg samples, gradually add BPW to the sample. Add a small portion of sterile BPW and mix to obtain a homogeneous suspension. Add the remainder of the BPW. Mix until a lump-free suspension is obtained.
- e. Incubate at $35 \pm 2^{\circ}\text{C}$ for 18-24 h.
- f. Proceed to Section 4.6 for use of the rapid screen or refer to Section 4.7 to continue the cultural analysis.

4.5.7 Raw Siluriformes (Fish) Products

Follow program requirements for preparing sample and sub-sample composites.

- a. Weigh the tissue into a sterile bag, sterile blender jar, or other sterile jar.
- b. Add the BPW. Stomach or blend, as required, for approximately two minutes or shake thoroughly.
- c. Incubate at $35 \pm 2^{\circ}\text{C}$ for 22-26 h.

**United States Department of Agriculture
Food Safety And Inspection Service, Office of Public Health Science**

MLG 4.10		Page 11 of 17
Title: Isolation And Identification of <i>Salmonella</i> From Meat, Poultry, Pasteurized Egg, and Siluriformes (Fish) Products and Carcass and Environmental Sponges		
Revision: .10	Replaces: .09	Effective: 01/02/19

- d. Proceed to Section 4.6 for use of the rapid screen or refer to Section 4.7 to continue the cultural analysis.

4.5.8 Fermented Products

Follow the procedure for RTE foods in Section 4.5.1 except:

- a. Blend/stomach the sample with 10 ± 0.2 g of sterilized calcium carbonate.
- b. Use buffered peptone water that contains 1 ml of a 1% aqueous solution of crystal violet per liter.

4.5.9 Dried Products (Breeding Mix, Dehydrated Sauce, Dried Soup Mix, and Dried Milk)

- a. Weigh the product into a sterile polypropylene bag, sterile blender jar, or other sterile jar.
- b. Add a small portion of the ambient temperature sterile BPW and mix to obtain a homogeneous suspension. Add the remainder of the BPW. Mix until a lump-free suspension is obtained.
- c. Incubate at $35 \pm 2^\circ\text{C}$ for 18-24 h.
- d. Proceed to Section 4.6 for use of the rapid screen or refer to Section 4.7 to continue the cultural analysis.

Note: Dried products such as soup mixes may require a sample/broth dilution greater than 1:10 because of physical difficulties encountered by absorption of broth by dehydrated product.

**United States Department of Agriculture
Food Safety And Inspection Service, Office of Public Health Science**

MLG 4.10	Page 12 of 17	
Title: Isolation And Identification of <i>Salmonella</i> From Meat, Poultry, Pasteurized Egg, and Siluriformes (Fish) Products and Carcass and Environmental Sponges		
Revision: .10	Replaces: .09	Effective: 01/02/19

4.5.10 Most Probable Numbers (MPN) Determination

Follow MPN instructions given in the specific project protocol or see MLG Appendix 2, Most Probable Number Procedure and Tables.

4.6 Rapid Screening Salmonella Test Procedure

Following incubation, perform the rapid screen using the current 3M™ Molecular Detection System User guide, or equivalent rapid screen technology for preparing reagents, performing the remainder of the assay, and reading the results.

- a. Samples that are rapid screen-negative will be reported as negative. All other samples will continue to cultural analysis as per MLG 4, Section 4.7. Alternatively, for samples with rapid screen results that are considered inconclusive, the laboratory may investigate. Based on the findings, the laboratory may:
 - repeat the rapid screen analysis from the lysate step or
 - prepare new rapid screen lysate tubes and repeat the analysis.
- b. In analytical runs where the positive control results are NOT positive, all samples are affected and an investigation shall be performed. Based on the findings the laboratory may:
 - repeat the rapid screen analysis from the lysate step
 - prepare new rapid screen lysate tubes and repeat the analysis or
 - analyze all of the samples culturally.

If circumstances (e.g. a power outage or equipment failure) do not allow testing using the rapid screen system, the laboratory shall, if possible, continue cultural analysis of all samples by proceeding with isolation and purification steps as per MLG 4, Section 4.7.

4.7 Selective Enrichment and Plating Media

- a. Transfer 0.5 ± 0.05 ml of sample into 10 ml TT broth (Hajna) broth and 0.1 ± 0.02 ml into 10 ml mRV broth.
- b. Incubate at $42 \pm 0.5^\circ\text{C}$ for 22-24 h or in a water bath at $42 \pm 0.5^\circ\text{C}$ for 18-24 h.

**United States Department of Agriculture
Food Safety And Inspection Service, Office of Public Health Science**

MLG 4.10	Page 13 of 17	
Title: Isolation And Identification of <i>Salmonella</i> From Meat, Poultry, Pasteurized Egg, and Siluriformes (Fish) Products and Carcass and Environmental Sponges		
Revision: .10	Replaces: .09	Effective: 01/02/19

- c. Carefully mix contents of tube by vortexing or equivalent means. Streak to BGS and DMLIA agar plates using a 10 μ l loopful of inoculum for each plate. Streak the entire agar plate with a single sample enrichment.
- d. Incubate at $35 \pm 2^{\circ}\text{C}$ for 18-24 h.
- e. Select typical colonies.

4.8 Examination of and Picking Colonies from Plating Media

4.8.1 Picking Colonies

- a. After the recommended incubation interval, examine the selective-differential agar plates and controls for the presence of colonies meeting the description for suspect *Salmonella* colonies. Pick well-isolated colonies.
 - BGS. Select colonies that are pink and opaque with a smooth appearance and entire edge surrounded by a red color in the medium. On very crowded plates, look for colonies that give a tan appearance against a green background.
 - DMLIA. Select purple colonies with (H_2S positive) or without (H_2S negative) black centers. Since *Salmonella* typically decarboxylate lysine and ferment neither lactose nor sucrose, the color of the medium reverts to purple.
- b. Pick at least one typical isolated colony from any of the plates. (NOTE: Before any sample is reported as *Salmonella* negative, pick at least three total typical colonies, if available. A representative of each typical colony type must be picked from each plate type before reporting the sample as *Salmonella* negative). Pick only from the surface and center of the colony. Avoid touching the agar because these highly selective media suppress growth of many organisms that may be viable.
- c. If there are typical colonies on a plate that are not well isolated, pick from the typical colonies and streak directly to a new set of selective agar plates. Alternatively, transfer typical colonies into a tube of TT broth (Hajna) or mRV broth and incubate overnight, then streak to selective agars.
- d. Incubate all plates for an additional 18-24 h at $35 \pm 2^{\circ}\text{C}$.

**United States Department of Agriculture
Food Safety And Inspection Service, Office of Public Health Science**

MLG 4.10	Page 14 of 17	
Title: Isolation And Identification of <i>Salmonella</i> From Meat, Poultry, Pasteurized Egg, and Siluriformes (Fish) Products and Carcass and Environmental Sponges		
Revision: .10	Replaces: .09	Effective: 01/02/19

- e. Reexamine initially negative plates and pick colonies as above. After 48-hour incubation, plates with no typical colonies may be discarded as negative. Plates with colonies undergoing confirmation testing should be stored at 2-8°C until testing is complete. If suspect *Salmonella* colonies do not confirm, reexamine the plates from which they were picked, and if appropriate, select colonies again for confirmation following Section 4.8.1.b.

4.8.2 Screening Media

- a. Inoculate TSI and LIA slants in tandem with a single pick from a colony by stabbing the butts and streaking the slants in one operation. If screw cap tubes are used, the caps must be loosened. Incubate at $35 \pm 2^\circ\text{C}$ for 24 ± 2 h.

Note: The same colony may be used to streak a plate used for subsequent biochemical testing described in Section 4.9.

Examine TSI and LIA slants as a set. Note the colors of butts and slants, blackening of the media, and for TSI slants presence of gas as indicated by gas pockets or cracking of the agar. Note also the appearance of the growth on the slants along the line of streak. A typical control on LIA should produce a purple butt with (H₂S positive) or without (H₂S negative) blackening of the media. A typical control on TSI should produce a yellow butt and red slant, with (H₂S positive) or without (H₂S negative) blackening of the media.

Discard, or re-streak for isolation, any sets that show "swarming" from the original site of inoculation. Discard sets that show a reddish slant in lysine iron agar. Isolates giving typical *Salmonella* sp. reactions and isolates that are suggestive but not typical of *Salmonella* sp. should be confirmed by a combination of biochemical and serological procedures. Refer to Table 2 for a summary of TSI-LIA reactions.

Note: TSI and LIA slants may be held at 2-8°C for up to 96 hours. Prior to any subsequent testing, fresh TSI and LIA slant(s) must be inoculated and incubated per instructions in a (above).

**United States Department of Agriculture
Food Safety And Inspection Service, Office of Public Health Science**

MLG 4.10	Page 15 of 17	
Title: Isolation And Identification of <i>Salmonella</i> From Meat, Poultry, Pasteurized Egg, and Siluriformes (Fish) Products and Carcass and Environmental Sponges		
Revision: .10	Replaces: .09	Effective: 01/02/19

- b. The motility testing in the last column of the table is optional. Refer to "Edwards and Ewing's Identification of Enterobacteriaceae" (Ewing, 1986) for additional information.
- c. Molecular serotyping or whole genome sequencing may be performed to further characterize isolates in lieu of serological testing with Somatic (O) Antigen and Flagellar (H) Antigen Agglutination Tests.

4.9 Biochemical Procedures

Commercially available biochemical test kits, including automated systems may be used for biochemical identification. Alternatively, use traditional methods of biochemical identification. Refer to AOAC Official Method 967.27 or "Edwards and Ewing's Identification of Enterobacteriaceae", 4th Edition, for biochemical reactions of *Enterobacteriaceae* and for fermentation media and test procedures.

For some biochemical test kits, i.e. the VITEK® 2 Compact System, streak a TSA + 5% sheep blood agar (SBA) plate from either the TSI slant, LIA slant, or a single colony pick from the plating media if inoculating TSI and LIA slants in tandem. Incubate 16-24 h at 35 ± 2°C. If using a UV positive control, perform a fluorescence check on the SBA plate for the positive control and presumptive positive samples following incubation. Use long wave UV light to examine plates for purity and evidence of cross contamination with the positive control. Only the positive control culture should fluoresce. If the presumptive positive sample SBA plates are pure and uncontaminated, perform the biochemical test.

Some commercial biochemical test systems may require streaking to other non-selective media prior to inoculation of their test kit. Follow manufacturer's instructions.

Note: Plates may be stored up to 96 hours at 2-8°C. Stored plates must be streaked to a new plate and incubated per instruction prior to use.

**United States Department of Agriculture
Food Safety And Inspection Service, Office of Public Health Science**

MLG 4.10	Page 16 of 17
Title: Isolation And Identification of <i>Salmonella</i> From Meat, Poultry, Pasteurized Egg, and Siluriformes (Fish) Products and Carcass and Environmental Sponges	
Revision: .10	Replaces: .09
Effective: 01/02/19	

Table 2. Potential *Salmonella* Reactions Requiring Biochemical Analysis. Additional culture work may be required and other factors should be considered before discarding any sample. Not all reactions are covered in this chart.

Triple Sugar Iron Agar			Lysine Iron Agar		O group and H antigen		Further Testing/ Disposal
Butt	Slant	H ₂ S	Butt	H ₂ S	O	H	
Y	R	+	P	+	+	+	B. & M. T.
Y	R	+	P	+	+	-	B. & M. T.
Y	R	-	P	-			B. & M. T.
Y	R	-	Y	-	+	+	B. & M. T.
Y	R	-	Y	-	-	-	B. & M. T.
Y	R	+	Y	+/-			B. & M. T.
Y	Y	-	Y or P	-			Discard
Y	Y	+	P	+			B. & M. T.
NC	NC						Discard

Y = Yellow (or A = acid reaction); R = Red (or K = alkaline reaction); P = Purple (or K = alkaline reaction);
 B. & M. T. = Perform biochemical testing, e.g., VITEK[®] 2 Compact System, and optional motility tests;
 NC = No change in color from uninoculated medium.

**United States Department of Agriculture
Food Safety And Inspection Service, Office of Public Health Science**

MLG 4.10		Page 17 of 17
Title: Isolation And Identification of <i>Salmonella</i> From Meat, Poultry, Pasteurized Egg, and Siluriformes (Fish) Products and Carcass and Environmental Sponges		
Revision: .10	Replaces: .09	Effective: 01/02/19

4.10 Culture Storage and Maintenance

For short-term (no longer than 3 months) storage, inoculate a nutrient agar slant, incubate at $35 \pm 2^{\circ}\text{C}$ overnight and then store at $2-8^{\circ}\text{C}$. For long-term storage, lyophilize cultures or freeze at $\leq -70^{\circ}\text{C}$ using cryo-beads, i.e. Cryostor™ or equivalent.

Maintain “working” *Salmonella* stock cultures on nutrient agar slants or equivalent. Transfer stocks monthly onto duplicate nutrient agar slants, incubate overnight at $35 \pm 2^{\circ}\text{C}$, and then maintain them at $2-8^{\circ}\text{C}$. Use one of the slants as the working culture. Use the other slant for sub-culturing to reduce the opportunity for contamination.

4.11 Selected References

Bailey, J. S., J. Y. Chiu, N. A. Cox, and R. W. Johnston. 1988. Improved selective procedure for detection of salmonellae from poultry and sausage products. *J. Food Prot.* **51**:391-396.

Centers for Disease Control and Prevention and National Institutes of Health (CDC/NIH). 2007. *BioSafety in Microbiological and Biomedical Laboratories*, 5th ed. or current. <http://www.cdc.gov/od/ohs/biosfty/bmb15/bmb15toc.htm>

Horowitz, William. (ed.). 2000. *Official methods of analysis of AOAC International*, 17th Edition. AOAC International Inc., Gaithersburg, MD 20877.

Ewing, W. H. 1986. *Edwards and Ewing's Identification of Enterobacteriaceae*, 4th Edition. Elsevier Science Publishing Co., Inc., New York.

Vassiliadis, P., D. Trichopoulos, A. Kalandidi, and E. Xirouchaki. 1978. Isolation of salmonellae from sewage with a new procedure of enrichment. *J. Appl. Bacteriol.* **66**:523-528.

Vassiliadis, P. 1983. The Rappaport-Vassiliadis (RV) enrichment medium for the isolation of *Salmonellas*: an overview. *J. Appl. Bacteriol.* **54**:69-76.

3M® Molecular Detection System User Guide